

Effectiveness of physical activity interventions in older adults with frailty or prefrailty: a systematic review and meta-analysis

Megan Racey PhD, Muhammad Usman Ali MD MSc, Diana Sherifali RN PhD, Donna Fitzpatrick-Lewis MSW, Ruth Lewis BA, Milos Jovkovic MSc, Danielle R. Bouchard PhD, Anik Giguère PhD, Jayna Holroyd-Leduc MD, Ada Tang PhD, Leah Gramlich MD, Heather Keller RD PhD, Jeanette Prorok PhD, Perry Kim PhD, Amanda Lorbergs PhD, John Muscedere MD; for the Canadian Frailty Network

See related research by Racey and colleagues at www.cmajopen.ca/lookup/doi/10.9778/cmajo.20200248

Abstract

Background: Physical activity is known to prevent frailty and reduce its consequences; however, it remains unclear which interventions are optimal for older adults with frailty. We conducted a systematic review and meta-analysis to identify effective physical activity interventions in improving outcomes related to frailty.

Methods: We searched MEDLINE, Embase, the Cochrane Central Register of Controlled Trials and CINAHL (inception to July 2019) for English-language randomized controlled trials (RCTs) of physical activity interventions in adults aged 65 years or more who were prefrail or frail; we included observational and cohort studies when there were no RCT data. Outcomes of interest were frailty, mobility, physical function, cognitive function, use of health care services and quality of life. After data extraction, we assessed the risk of bias using the Cochrane Collaboration risk of bias tool for RCTs and the Newcastle–Ottawa Scale for observational studies, rated the certainty of evidence with the Grading of Recommendation, Assessment, Development and Evaluations (GRADE) approach, and assessed statistical and methodologic heterogeneity.

Results: We identified 26 studies (24 RCTs [1 of which did not have any relevant outcomes for extraction] and 2 observational studies) involving 8022 prefrail or frail older adults. Nine studies had low risk of bias, 2 had high risk of bias, and for 13 the risk of bias was unclear. The trials included mixed (aerobic and muscle-strengthening) ($n = 13$), muscle-strengthening ($n = 8$), mobilization and rehabilitation ($n = 4$) or aerobic ($n = 1$) activities. Significant effects were found for mobility (standardized mean difference [SMD] 0.60, 95% confidence interval [CI] 0.37 to 0.83), activities of daily living (SMD 0.50, 95% CI 0.15 to 0.84), cognitive function (SMD 0.35, 95% CI 0.09 to 0.61), quality of life (SMD 0.60, 95% CI 0.13 to 1.07) and frailty (SMD -1.29 , 95% CI -2.22 to -0.36 ; risk ratio 0.58, 95% CI 0.36 to 0.93), with moderate certainty of evidence.

Interpretation: There is low- to moderate-level evidence that various physical activity interventions are beneficial for prefrail and frail older adults. Studies need to better define frailty to ensure the identification and implementation of such interventions into clinical practice. **PROSPERO registration:** CRD42020144556

Frailty is a leading contributor to functional decline and premature mortality in older adults.¹ More than 1.5 million Canadians are currently diagnosed as medically frail, and this number is expected to rise to more than 2 million in the next 10 years.² Frailty is a syndrome resulting from multiple factors and impairments that can reduce a person's functional ability. One of the major components of frailty is decline in physiologic domains such as loss of muscle mass and strength, flexibility, balance, coordination and performance.³ Older adults with frailty are at increased risk for falls, decline in mobility, hospital admission and death,^{4,5} which results in high consumption of health care resources,⁶ an increased burden on caregivers and adverse health outcomes.⁷

With the increase in older adults with frailty and the social and economic impact of this syndrome, research has focused on ways to prevent, delay and treat frailty, but proper identification and measurement of frailty is necessary to measure

Competing interests: Jeanette Prorok, Perry Kim, Amanda Lorbergs and John Muscedere are salaried employees of the Canadian Frailty Network. No other competing interests were declared.

This article has been peer reviewed.

Correspondence to: Diana Sherifali, dsherif@mcmaster.ca; Megan Racey, racey@mcmaster.ca

CMAJ Open 2021. DOI:10.9778/cmajo.20200222

these changes and optimize care and treatment.² Evidence shows a link between regular physical activity and improved muscle strength, aerobic capacity and balance in older adults in general⁸ and in those with frailty.^{9–11} However, much of this research has been conducted in populations in which frailty has not been clearly defined or measured. This has led to a gap in the literature: recommended physical activity^{12,13} may be too advanced or too intense for a frailer population, putting them at risk for falls and injuries.

Previous reviews looking at physical activity interventions for frail older adults have given mixed results,^{4,9,14–16} so it is still unclear what the best interventions are to support these people, especially in a population that has been clearly identified as frail. Such a defined population is important to support the creation of guidelines to enable evidence-based clinical practice.

We conducted a systematic review and meta-analysis focused on physical activity interventions in older adults with prefrailty or frailty (as identified by a tool or assessment) to assess the effectiveness of the interventions in improving outcomes including frailty, mobility, physical function (e.g., activities of daily living [ADLs], fatigue level, falls), cognitive function, use of health care services (e.g., hospital admissions) and quality of life. The results of this review, along with those of a methodologically similar review focused on nutrition and combined nutrition plus physical activity interventions,¹⁷ will provide the scientific evidence for the clinical practice guidelines of the Canadian Frailty Network.

Methods

This systematic review and meta-analysis followed the Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) guidelines.¹⁸ The protocol was registered in the International Prospective Register of Systematic Reviews (PROSPERO-CRD42020144556). There were minor variations from the published protocol. These included application of the English-only language restriction on screening rather than as part of the search, and reporting on only the outcomes listed in the protocol that were ranked critical based on the voting of a guideline panel committee. Our methods followed the Cochrane Handbook for Systematic Reviews of Interventions.¹⁹

Key question

What is the effectiveness of physical activity interventions in older adults (aged ≥ 65 yr) with frailty or prefrailty on clinical outcomes, outcomes that are important to the patient and outcomes related to use of health care services?

Search strategy

We developed the search terms, databases and strategy in consultation with a research librarian; they were informed by previous systematic reviews^{14–16} (Appendix 1, available at www.cmajopen.ca/content/9/3/E728/suppl/DC1). We searched MEDLINE (1948 to July 2019), Embase (1974 to July 2019), the Cochrane Central Register of Controlled

Trials (2019, Issue 6) and CINAHL (1937 to July 2019) and manually searched reference lists of relevant reviews and included studies not captured in our search. The search results were deduplicated, and citations were uploaded to a secure Internet-based platform for screening (DistillerSR, Evidence Partners).

Eligibility criteria

The PICO (population, intervention, comparator, outcomes) inclusion and exclusion criteria are provided in Box 1.

We included English-language peer-reviewed studies primarily from randomized controlled trials (RCTs); we included observational and cohort studies only when there were no RCT data. There was no exclusion based on intervention or participant setting.

Outcomes of interest were selected by an interdisciplinary steering committee (L.G., H.K., J.H.-L., A.G., A.T., D.R.B., J.M.) through a voting process that involved gathering a comprehensive list of outcomes from clinical and research expertise as well as existing relevant systematic reviews. The committee identified any missing outcomes and then anonymously rated the importance of the outcomes on a scale from 1 to 9 (1–3: not important; 4–6: important; 7–9: critical). Authors not involved in the ranking process (D.S., D.F.-L., M.R.) averaged the scores for each outcome and provided the list of outcomes to the committee for final discussion and agreement.

Box 1: Population, intervention, comparator, outcomes (PICO) eligibility criteria

Population

- Adults aged ≥ 65 yr clearly identified as prefrail or frail with any frailty assessment tool, assessment of frailty or other explicit author-established criteria (including studies with subanalyses involving a portion of prefrail or frail participants)
- To make this review relevant to the general frail population, studies that targeted clinical populations, such as patients with obesity or cancer, were excluded

Intervention

- Physical activity interventions

Comparator

- True control group defined as having received usual care, routine care or minimal contact that did not include any intervention or treatment group components
- Any head-to-head interventions were excluded

Outcomes

- Frailty (measured with a valid tool)
- Mobility (e.g., gait speed, Timed Up and Go test, Sit-to-Stand test, balance test and Short Physical Performance Battery)
- Physical function (e.g., activities of daily living, energy and fatigue levels, falls, fall rate, fall incidence)
- Cognitive function (e.g., Loewenstein Occupational Therapy Cognitive Assessment, Mini-Mental State Examination)
- Use of health care services (e.g., hospital admission, emergency department visits)
- Quality of life (measured with a standardized tool)

Study selection, data extraction and quality assessment

A series of team members were involved in the screening process, which resulted in different combinations of reviewers and team members. At all levels of screening, all articles required a minimum of 2 reviews. Multiple publications for the same primary intervention were merged; all data from all publications from a single study were extracted for relevant outcomes, but we describe the study and its characteristics only once in our review.

We developed, piloted and deployed standardized forms for data extraction. These forms were templated based on our experience and expertise as methodologists and included details such as outcome type, tool used, and intervention and control group data. The forms were then tested independently by 2 researchers to ensure there were no errors or misinterpretation. Two team members then completed full data extraction (unadjusted, intention-to-treat data) and an assessment of risk of bias using the Cochrane Collaboration risk of bias tool²⁰ for RCTs and the Newcastle–Ottawa Scale²¹ for observational studies. If interventions had multiple treatment arms, only the interventions that met our inclusion criteria were extracted. Conflicts were resolved by the lead researcher of this review (M.R.), and all data extraction was verified independently by the statistician (M.U.A.).

Data on harms or adverse events were extracted narratively.

Certainty of evidence

We independently evaluated the certainty of the body of evidence using the Grading of Recommendation, Assessment, Development and Evaluations (GRADE) approach²² with GRADEpro software. With this approach, the certainty of a body of evidence is rated as high, moderate, low or very low based on assessment of 5 conditions: methodologic quality; consistency across effect estimates and statistical heterogeneity; directness of the body of evidence to the populations, interventions, comparators or outcomes of interest; precision of results; and indications of reporting bias.

Statistical analysis

The physical activity components and intensities extracted from the search are shown in Appendix 2 (available at www.cmajopen.ca/content/9/3/E728/suppl/DC1). All data analyses were planned a priori. We used a meta-analysis to combine the results across all studies by outcome using the published data from included studies (full methods presented in Appendix 3, available at www.cmajopen.ca/content/9/3/E728/suppl/DC1).

We also conducted an analysis of the effectiveness of each physical activity category by outcome. For continuous outcomes, we used the change from baseline to immediately after the intervention (mean change score, standard deviation [SD] of mean change score) for both the intervention group and the control group to generate the summary measures of effect in the form of standardized mean difference (SMD).²³ We used SMD as a summary statistic because many studies in this systematic review assessed the same outcome measured in a

variety of ways (e.g., mobility measured as stair climb, balance test, gait speed, chair rise repetition, Sit-to-Stand test, Short Physical Performance Battery, gait speed, Timed Up and Go test). In this situation, it was necessary to standardize the results of the studies before they could be compared across studies or combined in a quantitative synthesis. The SMD-based effect sizes represent the magnitude of the intervention effect relative to the variability observed in a particular study. Therefore, studies for which the difference in mean change score was the same as the proportion of SD of mean change score have the same SMD, regardless of the actual scale or unit of measurement used to obtain the outcome measures.^{19,24} The SMD is interpreted based on its magnitude according to recommended Cohen *d* thresholds (~0.2 = small effect, ~0.5 = medium effect, ~0.8 = large effect).²⁵ For studies in which measure of variance was reported as confidence intervals (CIs), standard error or *p* values, we used Cochrane-recommended methods to convert these data to SD.¹⁹

We used a multilevel meta-analytical approach (where applicable) to account for statistical dependence, that is, dependency in effect sizes introduced by comparison of multiple intervention arms within a study to a common control group, or by multiple outcome measures or suboutcome measures of a primary outcome of interest within a study (such as Short Physical Performance Battery reported as gait speed, balance and chair stand test separately).²⁶ In such cases, we nested the correlated measures or effect sizes within studies first, by introducing a random effect to our grouping variables such as studies, outcome measures and intervention arms. This grouping variable, also known as a random intercept, told our model to assume different values (intercepts) for each grouping level. Specifically, we used 2 grouping variables: that on level 2 and that on level 3. We assumed that these grouping variables are nested, in the sense that several effect sizes on level 2 together make up a larger cluster on level 3. For pooling of performance measures, we adjusted the direction of effect to ensure consistency of desirable outcome responses.

For dichotomous outcomes, we used the number of events after the intervention to generate the summary measures of effect in the form of risk ratio (RR) using DerSimonian and Laird random-effects models with the Mantel–Haenszel method. We used the Cochran Q test ($\alpha = 0.05$) to detect statistical heterogeneity and the *I*² statistic to quantify the magnitude of statistical heterogeneity between studies, where an *I*² value of 30%–60% represents moderate heterogeneity and a value of 60%–90% represents substantial heterogeneity across studies.

We estimated the statistical heterogeneity *I*² statistic in the context of the multilevel meta-analytical approach.^{24,26} We estimated *I*²-level 2 to account for the amount of within-cluster heterogeneity (i.e., across effect sizes or multiple arms of same study), *I*²-level 3 to account for between-cluster heterogeneity (i.e., effect sizes across studies or subgroups of interest) and *I*²-total to represent the heterogeneity not attributable to sample error (sum of values at levels 2 and 3). We added *I*²-total to all forest plots for overall pooled effect estimates and summary effect size within subgroups.

We assessed publication bias using funnel plots²⁷ when there were at least 10 studies in the meta-analysis.

We performed all analyses using R software (metafor and dmetar packages) (R Foundation for Statistical Computing).

Ethics approval

As this study was solely literature based, it was not eligible for institutional ethics approval, and none was sought.

Results

From the 4668 citations identified, we assessed 218 full-text articles for eligibility and included 26 studies described in 34 publications in the qualitative synthesis (Figure 1); the studies were published from 1998 to 2019. Of the 26 studies, 24 were RCTs^{28–51} and 2 were observational studies;^{52,53} however, 1 of

the RCTs did not have any relevant outcomes for extraction.⁵¹ Therefore, we meta-analyzed 23 RCTs and their outcomes.

The 26 studies included in the qualitative analysis involved 8022 frail older participants aged 65 years or more (mean 69.8–85.4 yr); the proportion of women ranged from 50% to 100%. All of the included RCTs had fewer than 280 participants (median 88; quartiles 62, 88, 148, 1635), with the exception of 1, which had 1635 participants.³⁰

Characteristics of the included studies are presented in Table 1, and further demographic data from the studies can be found in Appendix 4 (available at www.cmajopen.ca/content/9/3/E728/suppl/DC1). Briefly, interventions were conducted in Asia, Europe and North America (all in the United States) with participants who were community-dwelling or living in care homes ($n = 12$), or had been admitted to hospital.

The tools used to measure and assess frailty included standardized tools such as Fried's frailty phenotype but also other assessments of mobility, physical dependency and ADL indices. Most interventions ($n = 13$) consisted of both aerobic and muscle-strengthening activities, known as the mixed (or multicomponent) category, followed by muscle-strengthening activities only ($n = 8$) (Table 2). The intensity of the interventions was mostly resistance or strength training ($n = 9$), followed by moderate intensity ($n = 7$). Activities were performed 1–2 or 3–4 times per week and were 30–60 minutes in duration. The interventions lasted on average 29 weeks, with only 4 studies having a duration of 9 months or more.

Adverse effects or harms that were directly related to the intervention were reported in 7 studies (5 mixed physical activity,^{36,43,45,52,53} 1 muscle-strengthening intervention³¹ and 1 mobilization or rehabilitation intervention⁴⁴), and no occurrences of any adverse effects were reported in 5. Most of the harms consisted of falls (occurring during the intervention protocol or physical activities); fractures, sprains, strains or other injuries; muscle ache and fatigue; and incidents from other pre-existing health problems, such as angina; there was no significant difference between the intervention and control groups. Events were successfully managed with adjustments to intervention protocols in most cases, and participant retention remained high (79%⁴⁴ to 92%³¹), but 3 participants withdrew from 2 studies.^{43,45}

Risk of bias and quality of studies

Using the Cochrane Collaboration risk of bias tool, we found that 9 studies had low risk of bias^{28,36,37,41–44,48,49} and 2 studies had high risk of bias;^{32,34} the risk of bias was unclear for 13 studies,^{29–31,33,35,38–40,45–47,50,51} mostly owing to unclear sequence generation, allocation concealment and blinding procedures (Table 3).

The certainty of evidence as assessed with the GRADE approach ranged from very low to moderate; it was moderate for most outcomes owing to downgrading for risk of bias (Table 4; Appendix 5, available at www.cmajopen.ca/content/9/3/E728/suppl/DC1).

We did not observe any significant asymmetry across funnel plots for publication bias (Appendix 6, available at www.cmajopen.ca/content/9/3/E728/suppl/DC1).




Figure 1: Flow diagram showing study selection. Note: RCT = randomized controlled trial.

Table 1 (part 1 of 3): Characteristics of the included studies

Study; country	No. of participants (F/M, %)	Age, mean ± SD, yr	Assessment tool or measure*	Frailty level, no. of participants†	Design; duration‡	Intervention category; intensity; frequency; duration	Control	Delivery of intervention	Outcome(s)
Binder et al., ⁵⁴⁻⁵⁶ 2005, US	119 (52/48§)	O: 83 ± 4 I: 83 ± 4 C: 83 ± 4	Physical Performance Test¶	I: mild to moderate frailty 66 C: mild to moderate frailty 49	RCT; 9 mo	Mixed; moderate intensity; 3x/wk; 30 min (aerobic component)	Home exercises (primarily flexibility), monthly exercise classes	Exercise physiology technician	Frailty
Brown et al., ⁴⁶ 2000, US	87 (57/43)	I: 83 ± 4 C: 83 ± 4	Physical Performance Test	I: frail 48 C: frail 39	RCT; 3 mo	Mobility/rehabilitation; light intensity; 3x/wk; duration NR	Home exercises (primarily flexibility), monthly exercise classes	NR	Frailty, mobility, physical function (including falls and quality of life)
Chandler et al., ⁴⁸ 1998, US	100 (50/50)	O: 77.6 ± 7.6 I: 77.5 ± 7.1 C: 77.7 ± 7.8	Inability to descend stairs step over step without holding railing	I: frail 50 C: frail 50	CCT; 10 wk	Muscle-strengthening; resistance/strength training; 3x/wk; duration NR	Usual care	Physiotherapist	Frailty, physical function (including falls and quality of life)
Chen et al., ⁵¹ 2010, Taiwan	69 (53/47)	O: 75.40 ± 6.70	Barthel Index (ADLs)	I: transitional frail 38 C: transitional frail 31	RCT; 8 mo	Mobility/rehabilitation; light intensity; 3x/wk; 70 min	Usual care	Certified yoga instructor	Frailty
Chen et al., ²⁸ 2020, China	70 (65/35§)	I: 76.97 ± 5.19 C: 75.27 ± 5.98	Fried's frailty phenotype	I: prefrail 33 C: prefrail 33	RCT; 9 wk	Muscle-strengthening; resistance/strength training; 3x/wk; 45-60 min	Usual care	Physiotherapist or trained community centre worker	Frailty, mobility
Clegg et al., ³¹ 2014, UK	84 (71/29)	O: 79 ± 9.2 I: 79.4 ± 7.9 C: 78.0 ± 10.5	Edmonton Frail Scale	Mean Edmonton Frail Scale score ± SD: I: 7.8 ± 2.4 C: 8.3 ± 2.7	RCT; 12 wk	Muscle-strengthening; resistance/strength training; 5x/wk; < 15 min	Usual care	Physiotherapist	Frailty, mobility
Daniel, ⁴⁰ 2012, US	23 (61/39)	O: 77 ± 5.3 I: 78.13 ± 5.5, 80 ± 3.37 C: 72.6 ± 4.6	Fried's frailty phenotype	I: prefrail 8, 8 C: prefrail 7	RCT; 15 wk	Mixed; intensity unknown; 3x/wk; 45 min	Usual care	Certified fitness professional	Frailty, mobility
de Jong et al., ^{45,57,58} 2000, the Netherlands	217 (70/30)	O: 78.7 ± 5.6 I: 76.7 ± 4.4 C: 79.3 ± 6.6	Required health care service (e.g., home care or meal delivery)	I: frail 39 C: frail 37	RCT; 17 wk	Mixed; moderate intensity; 2x/wk; 45 min	Social program	Teacher (researcher supervised)	Frailty, mobility, physical function (including falls and quality of life)
Faber et al., ⁴² 2006, the Netherlands	278 (79/21§)	I: 85.4 ± 5.9, 84.4 ± 6.4 C: 84.9 ± 5.9	Fried's frailty phenotype¶	I: prefrail 25, 33; frail 36, 43; not frail 4, 4 C: prefrail 47; frail 36; not frail 7	RCT; 20 wk	Muscle-strengthening; moderate intensity; 1x/wk for 4 wk, then 2x/wk for 16 wk; 90 min	Usual care	Instructor-led group sessions	Frailty, mobility, physical function (including falls and quality of life)
Gill et al., ^{44,59} 2002, US	188 (80/20)	O: 83 I: 82.8 ± 5.0 C: 83.5 ± 5.2	Moderately frail if required > 10 s to perform rapid gait test or could not stand up from seated position in hardback chair with arms folded; severely frail if met both criteria	I: moderately frail 60, severely frail 34 C: moderately frail 56, severely frail 38	RCT; 7 mo	Mobility/rehabilitation; intensity unknown; 3x/wk; duration NR	Individualized monthly health education sessions	Physiotherapist	Frailty, mobility, cognitive function, physical function (including falls and quality of life), use of health care services

Table 1 (part 2 of 3): Characteristics of the included studies

Study; country	No. of participants (F/M, %)	Age, mean \pm SD, yr	Assessment tool or measure*	Frailty level, no. of participants†	Design; duration‡	Intervention category; intensity; frequency; duration	Control	Delivery of intervention	Outcome(s)
Giné-Garriga et al., ⁴¹ 2010, Spain	51 (61/39§)	I: 83.9 \pm 2.8 C: 84.1 \pm 3.0	> 10 s to perform rapid gait test, could not stand up 5 times from seated position in hardback chair with arms folded, or Fried's exhaustion criteria	I: frail 22 C: frail 19	RCT; 12 wk	Mixed; resistance/strength training; 2 \times /wk; 45 min	Usual care, social meetings	Researcher	Frailty, mobility, physical function (including falls and quality of life)
Giné-Garriga et al., ³⁹ 2013, Spain	51 (61/39§)	I: 83.9 \pm 2.8 C: 84.1 \pm 3	Fried's frailty phenotype	I: frail 22 C: frail 19	RCT; 12 wk	Mixed; resistance/strength training; 2 \times /wk; duration NR	Usual care, social meetings	NR	Frailty, physical function (including falls and quality of life)
Kuo et al., ⁴⁹ 2018, Taiwan	41 (83/17§)	O: 69.77 I: 68.93 \pm 3.81 C: 70.38 \pm 5.22	Comprehensive geriatric assessment	I: frail 15 C: frail 21	RCT; 8 wk	Aerobic; moderate intensity; 2 \times /wk; 30 min	Usual care	Researcher	Frailty, mobility, cognitive function, physical function (including falls and quality of life)
Kwon et al., ³⁷ 2015, Japan	89 (100/0††)	O: 76.8 I: 77.0 \pm 4.2 C: 76.9 \pm 3.9	Fried's frailty phenotype¶	I: prefrail 28 C: prefrail 31	RCT; 12 wk	Mixed; intensity unknown; 1 \times /wk; 60 min	General health education sessions	Certified health fitness trainer	Frailty, mobility, physical function (including falls and quality of life)
Liu et al., ³² 2017, Hong Kong	85 (94/6††)	O: 79.32 \pm 7.72 I: 79.72 \pm 7.95, 77.03 \pm 7.07‡‡ C: 81.90 \pm 7.69	Fried's frailty phenotype	I: prefrail 5, frail 24 C: prefrail 6, frail 15	RCT; 16 wk	Mixed; intensity unknown; 1 \times /wk at centre, 3 \times /wk at home; 45–60 min at centre, 30–45 min at home	Centre-based health talks on management of various health issues except fatigue	Physiotherapist and exercise instructor	Frailty, mobility, physical function (including falls and quality of life)
Liu et al., ³⁰ 2018, US	1635 (66/34§)	I: 78.6 \pm 5.2 C: 79.1 \pm 5.3	Study of Osteoporotic Fractures index	NR**	RCT; 2 yr	Mixed; moderate intensity; 3–4 \times /wk; 50 min	Health education workshops, instructor-led program of gentle upper extremity stretching or flexibility exercises	NR	Frailty, mobility
Losa-Reyna et al., ²⁹ 2019, Spain	30 (75/25§)	O: 84.2 \pm 4.5 I: 84.0 \pm 4.7 C: 84.4 \pm 4.6	Fried's frailty phenotype	O: prefrail 7, frail 13	CCT; 6 wk	Mixed; high/strenuous intensity; 2 \times /wk; 45 min	Usual care	NR	Frailty, mobility, physical function (including falls and quality of life)
Ng et al., ^{36,60,61} 2015, Singapore	246 (56/44††)	O: 70.0 \pm 4.7 I: 69.7 \pm 4.23, 69.7 \pm 4.31, 70.3 \pm 5.25, 70.4 \pm 4.74‡‡ C: 70.1 \pm 5.02	Cardiovascular Health Study criteria	I: prefrail 29, frail 19 C: prefrail 43, frail 7	RCT; 24 wk	Mixed; resistance/strength training; 2 \times /wk; 90 min	Standard care	Qualified trainer	Frailty, mobility, cognitive function, physical function (including falls and quality of life), use of health care services
Takatori et al., ³⁴ 2016, Japan	266 (100/0)	O: 75 \pm 5 I: 74.6 \pm 5.1 C: 75.9 \pm 6.0	Kihon Checklist	I: frail 148 C: frail 118	CCT; 6 mo	Muscle-strengthening; intensity unknown; 3 \times /wk; 5 min	Maintained or improved activity in daily life, lectures on health	Physiotherapist, then self-guided	Frailty, mobility, physical function (including falls and quality of life)

Table 1 (part 3 of 3): Characteristics of the included studies

Study; country	No. of participants (F/M, %)	Age, mean ± SD, yr	Assessment tool or measure*	Frailty level, no. of participants†	Design; duration‡	Intervention category; intensity; frequency; duration	Control	Delivery of intervention	Outcome(s)
Tarazona-Santabalbina et al., ³⁵ 2016, Spain	100 (54/46)	I: 79.7 ± 3.6 C: 80.3 ± 3.7	Fried's frailty phenotype	I: frail 51 C: frail 49	RCT; 24 wk	Mixed; moderate intensity; 5x/wk; 65–70 min	Usual care	Physiotherapist and nurse	Frailty, mobility, cognitive function, physical function (including falls and quality of life), use of health care services
Tieland et al., ³⁸ 2015, the Netherlands	127 (61/29)	O: 79.0 ± 0.7 I: 78.4 ± 1.0 C: 79.5 ± 1.0	Fried's frailty phenotype	I: prefrail 84%, frail 16% C: prefrail 72%, frail 28%	CCT; 24 wk	Muscle-strengthening; resistance/strength training; 2x/wk; duration NR	Usual care	Personal supervision	Frailty, mobility, physical function (including falls and quality of life)
Tsang et al., ⁵⁰ 2013, China	134 (75/25§)	I: 83.33 ± 6.30 C: 84.85 ± 6.03	62-item frailty index	I: frail 61 C: frail 55	RCT; 12 wk	Mobility/rehabilitation; light intensity; frequency NR; 60 min	Newspaper reading and discussion activity	Certified qigong instructor	Frailty, mobility, cognitive function
Yamada et al., ⁵³ 2012, Japan	610 (77/23)	I: 79.7 ± 6.3 C: 80.3 ± 6.6	Frailty checklist of Japan	Mean total score on frailty checklist ± SD: I: 7.41 ± 3.98 C: 7.34 ± 4.27	Prospective cohort study; 16 wk	Mixed; moderate intensity; 1x/wk; 90 min	Screening evaluation	Physiotherapist	Frailty
Yamada et al., ⁵² 2017, Japan	3240 (82/18)	I: 77.1 ± 6.4 C: 77.2 ± 6.9	Kihon Checklist	I: prefrail 591, frail 420, robust 609 C: prefrail 591, frail 420, robust 609	Prospective cohort study; 4 yr	Mixed; light intensity; 1x/wk; 60 min	Matched on all baseline characteristics (e.g., age, gender, body mass index) and each item on Kihon Checklist	Volunteer staff	Frailty
Yarasheski et al., ⁴⁷ 1999, US	17 (71/29)	I: female 82 ± 2, male 82 ± 1 C: female 82 ± 2, male: 82§§	Physical Performance Test, self-reported ADLs	I: mild to moderate frailty 12 C: mild to moderate frailty 5	RCT; 9 mo	Muscle-strengthening; resistance/strength training; 3x/wk; duration NR	Home exercises (primarily flexibility), monthly exercise sessions	NR	Frailty, physical function (including falls and quality of life)
Yoon et al., ³³ 2018, South Korea	65 (70/30§)	O: 73.94 ± 4.27 I: 73.82 ± 4.37 C: 74.03 ± 4.27	Cardiovascular Health Study criteria	Mean frailty criteria ± SD: I: 1.63 ± 0.90 C: 1.37 ± 0.56	RCT; 16 wk	Muscle-strengthening; resistance/strength training; 3x/wk; 60 min	Usual care, dynamic stretching	Exercise instructor	Frailty, mobility, cognitive function, physical function (including falls and quality of life)

Note: ADL = activity of daily living, C = control, CCT = clinical controlled trial, F = female, I = intervention, M = male, NR = not reported, O = overall, RCT = randomized controlled trial, SD = standard deviation.
 *Fried's frailty criteria were derived from the Cardiovascular Health Study (CHS) and as such can be considered the same as the CHS criteria; however, we report here how the assessment tool was described by the study authors.
 †Total nonfrail, prefrail and frail for entire study population by intervention and control groups (may include multiple treatment arms combined).
 ‡Not including follow-up, if applicable.
 §Values based on reported baseline, which may not equal the number randomly allocated but, rather, the number who completed the intervention.
 ¶Authors indicated that the assessment tool was modified.
 **Authors indicated that most participants were frail; however, the number of frail participants was unclear.
 ††The *n* from the most relevant intervention/treatment arms was used to calculate proportions.
 ‡‡Only 1 intervention group met the inclusion criteria for this review.
 §§Only 1 participant.

Benefits of treatment

The meta-analysis included an examination of the between-group impact of all physical activity RCTs together and subgroup analyses based on the intervention category (aerobic [*n* = 1], muscle-strengthening [*n* = 8], mobilization or rehabilitation [*n* = 4] and mixed [*n* = 13]). Owing to hetero-

geneity, we analyzed the physical outcomes separately based on the individual measures of ADLs, falls and fatigue level. Forest plots for meta-analyzed outcomes for all interventions and by intervention category can be found in Appendix 7 (available at www.cmajopen.ca/content/9/3/E728/suppl/DC1).

Table 2: Category, intensity, frequency, duration and delivery of included studies

Characteristic	No. (%) of studies <i>n</i> = 26
Physical activity category	
Aerobic	1 (4)
Mixed	13 (50)
Mobilization/rehabilitation	4 (15)
Muscle strengthening	8 (31)
Intervention intensity	
High/strenuous	1 (4)
Moderate	7 (27)
Light	4 (15)
Resistance/strength training	9 (35)
Unknown	5 (19)
Frequency of intervention	
1–2 times per week	12 (46)
3–4 times per week	11 (42)
> 4 times per week	2 (8)
Not reported	1 (4)
Duration of intervention session, min	
< 15	2 (8)
30–60	13 (50)
> 60	5 (19)
Not reported	6 (23)
Delivery agent	
Physiotherapist	9 (35)
Fitness instructor/trainer	7 (27)
Researcher	3 (12)
Other	2 (8)
Not reported	5 (19)

Overall

Overall, the RCTs had significant pooled effect estimates on measures of mobility (SMD 0.60, 95% CI 0.37 to 0.83) (Figure 2), ADLs (SMD 0.50, 95% CI 0.15 to 0.84) (Figure 3A), cognitive function (SMD 0.35, 95% CI 0.09 to 0.61) (Figure 3B), quality of life (SMD 0.60, 95% CI 0.13 to 1.07) (Figure 4A) and frailty (SMD -1.29, 95% CI -2.22 to -0.36; RR 0.58, 95% CI 0.36 to 0.93) (Figures 4B and 4C, Table 4; Appendix 7), with moderate certainty of evidence (Appendix 5, Supplemental Table S3). These effects were large for frailty, medium for mobility, ADLs and quality of life, and small for cognitive function. There were no significant effects on measures of falls or fatigue level (Appendix 7).

Aerobic interventions

The only aerobic intervention, which was a prescribed stepper walking program,⁴⁹ had a medium but significant effect estimate on mobility (SMD 0.71, 95% CI 0.23 to 1.20) (Table 4;

Appendix 7), with low certainty of evidence (Appendix 5, Supplemental Table S4). The mean age of participants was 69.8 years, and 83% were women.

Muscle-strengthening interventions

The muscle-strengthening RCTs had significant pooled effect estimates on measures of mobility (SMD 0.57, 95% CI 0.08 to 1.06) and frailty (RR 0.21, 95% CI 0.10 to 0.43) (Table 4; Appendix 7), with moderate certainty of evidence (Appendix 5, Supplemental Table S5), and a small but significant effect on the measure of cognitive function (SMD 0.45, 95% CI 0.19 to 1.72) (Table 4; Appendix 7), with very low certainty of evidence (Appendix 5, Supplemental Table S5). The mean age of participants ranged from 74 to 85 years; 70% were women.

Mobilization and rehabilitation interventions

One of the 4 mobilization or rehabilitation RCTs did not have any relevant outcomes for extraction.⁵¹ The 3 remaining RCTs had small but significant pooled effect estimates on measures of mobility (SMD 0.29, 95% CI 0.17 to 0.42) and ADLs (SMD 0.48, 95% CI 0.28 to 0.67) (Table 4; Appendix 7), with moderate certainty of evidence (Appendix 5, Supplemental Table S6). The mean age of participants ranged from 75.4 to 84.9 years; 70% were women.

Mixed interventions

The RCTs of mixed interventions had significant pooled effect estimates on measures of mobility (SMD 0.75, 95% CI 0.40 to 1.10), ADLs (SMD 0.64, 95% CI 0.004 to 1.27), cognitive function (SMD 0.62, 95% CI 0.12 to 1.11), quality of life (SMD 0.68, 95% CI 0.16 to 1.21) and frailty (SMD -1.57, 95% CI -2.57 to -0.57; RR 0.72, 95% CI 0.63 to 0.83) (Table 4; Appendix 7). These effects were large for frailty, with moderate certainty of evidence; medium for mobility, cognitive function and quality of life, with moderate certainty of evidence; and medium for ADLs, with low certainty of evidence (Appendix 5, Supplemental Table S7). The mean age of participants ranged from 75.9 to 84.1 years; 69% were women.

Both observational studies were prospective cohorts with mixed physical activity.^{52,53} Only 1 contained data for the outcomes selected in this review.⁵³ Evidence with low certainty indicates that there is a reduced risk of frailty with higher physical activity levels compared to lower physical activity levels (Appendix 8, available at www.cmajopen.ca/content/9/3/E728/suppl/DC1).

Interpretation

Our review showed a significant benefit of physical activity interventions of various types on certain outcomes including mobility, ADLs, cognitive function, quality of life and frailty when compared to control groups in frail adults aged 65 years or more. The effect sizes ranged from small to large, with low to moderate certainty of evidence. When we looked at all physical activity interventions together, there was a

Table 3: Risk of bias for the included studies

Study	Type of bias					
	Sequence generation	Allocation concealment	Blinding of outcome assessment	Incomplete outcome data	Selective reporting	Other
Chen et al., ²⁸ 2020	L	L	L	L	L	U
Losa-Reyna et al., ²⁹ 2019	U	U	U	H	L	L
Liu et al., ³⁰ 2018	L	U	U	L	L	H
Clegg et al., ³¹ 2014	U	U	H	U	L	U
Liu et al., ³² 2017	H	H	U	L	H	H
Yoon et al., ³³ 2018	U	U	U	H	L	L
Takatori et al., ³⁴ 2016	H	H	L	H	L	L
Tarazona-Santabalbina et al., ³⁵ 2016	U	U	L	L	L	U
Ng et al., ^{36,60,61} 2015	L	L	L	L	L	L
Kwon et al., ³⁷ 2015	L	L	L	L	L	L
Tieland et al., ³⁸ 2015	U	U	U	H	L	U
Giné-Garriga et al., ³⁹ 2013	L	U	H	H	L	L
Daniel, ⁴⁰ 2012	U	U	U	L	L	H
Giné-Garriga et al., ⁴¹ 2010	L	U	L	H	L	L
Faber et al., ⁴² 2006	L	U	L	L	L	H
Binder et al., ^{43,54-56} 2005	L	U	L	L	L	L
Gill et al., ^{44,59} 2002	L	U	L	U	L	U
de Jong et al., ^{45,57,58} 2000	U	U	U	U	L	H
Brown et al., ⁴⁶ 2000	U	U	U	L	L	U
Yarasheski et al., ⁴⁷ 1999	U	U	U	U	L	U
Chandler et al., ⁴⁸ 1998	L	U	L	L	L	L
Kuo et al., ⁴⁹ 2018	L	U	L	L	L	U
Tsang et al., ⁵⁰ 2013	U	U	U	L	L	H
Chen et al., ⁵¹ 2010	U	U	U	L	L	U

Note: H = high risk of bias, L = low risk of bias U = risk of bias unclear.

large effect on frailty, a medium effect on quality of life, ADLs and mobility, and a small effect on cognitive function.

Previous systematic reviews and meta-analyses have been conducted to evaluate various physical activity interventions and their outcomes related to frailty prevention, progression and reversal, but few have used such an explicit inclusion criterion for prefrailty or frailty, or combined individual measurements of outcomes (such as muscle strength and gait speed) into overall effect estimates (such as physical performance) to allow for a more robust presentation of the results.^{9-11,62-67} However, frailty itself was measured infrequently, both at baseline and after the intervention. The authors of only 4 of the 23 studies that we meta-analyzed reported frailty outcomes after the intervention,^{29,33,35,36} 3 of which were mixed physical activity interventions. When compared to control, these interventions showed a large effect size, with moderate certainty of evidence, similar to other reviews;^{5,63,66,68,69} however, the

GRADE approach was not used in any of those reviews, and only 2 included a meta-analysis of outcomes.^{63,69}

Mobility-related outcomes (gait speed, Timed Up and Go test, Sit-to-Stand test, Short Physical Performance Battery, balance test and Physical Performance Test) and ADLs (measured with valid tools such as the Barthel Index, Instrumental Activities of Daily Living Scale, Instrumental Activities of Daily Living instrument and Groningen Activity Restriction Scale) were significantly improved by almost all physical activity intervention types. Mobility was the most common outcome reported among the studies in our review (19/26). The significant pooled effect estimates were mostly the result of the mixed ($n = 9$) and muscle-strengthening ($n = 6$) interventions. Mixed training has been shown to be effective at managing frailty in older adults, with improvements in mobility-related outcomes such as muscle strength and balance, and reductions in disability and falls.^{15,62,63,66,68,69} Previous reviews have also

Table 4 (part 1 of 2): Physical activity intervention effect estimates for outcomes and certainty of evidence, for all interventions combined and by intervention category

Category; outcome*†	No. of studies	No. of participants	SMD‡ (95% CI)§	GRADE rating
All interventions combined				
Mobility	19	1724	0.60 (0.37 to 0.83)	Moderate; downgraded for risk of bias
Activities of daily living	9	910	0.50 (0.15 to 0.84)	Moderate; downgraded for risk of bias
Cognitive function	5	377	0.35 (0.09 to 0.61)	Moderate; downgraded for risk of bias
Quality of life	6	500	0.60 (0.13 to 1.07)	Moderate; downgraded for risk of bias
Frailty	4	244	-1.29 (-2.22 to -0.36)	Moderate; downgraded for risk of bias
	4	1538	RR 0.58 (0.36 to 0.93)¶	Moderate; downgraded for risk of bias
Falls	7	724	RR 0.80 (0.51 to 1.26)¶	Very low; downgraded for risk of bias, inconsistency and imprecision
Fatigue level	3	184	-0.27 (-0.65 to 0.12)	Low; downgraded for risk of bias and imprecision
Aerobic interventions				
Mobility	1	36	0.71 (0.23 to 1.20)	Low; downgraded for risk of bias and imprecision
Activities of daily living	1	36	0.46 (-0.03 to 0.94)	Very low; downgraded for risk of bias and imprecision
Cognitive function	1	36	0.15 (-0.50 to 0.80)	Very low; downgraded for risk of bias and imprecision
Fatigue level	1	36	-0.39 (-0.87 to 0.09)	Very low; downgraded for risk of bias and imprecision
Muscle-strengthening interventions				
Mobility	6	722	0.57 (0.08 to 1.06)	Moderate; downgraded for risk of bias
Activities of daily living	2	278	0.16 (-0.05 to 0.37)	Low; downgraded for risk of bias and imprecision
Cognitive function	1	45	0.45 (0.19 to 0.72)	Very low; downgraded for risk of bias and imprecision
Quality of life	1	70	0.15 (-0.33 to 0.63)	Very low; downgraded for risk of bias and imprecision
Frailty	1	45	-0.20 (-0.79 to 0.39)	Very low; downgraded for risk of bias and imprecision
	1	66	RR 0.21 (0.10 to 0.43)¶	Moderate; downgraded for imprecision
Falls	2	442	RR 0.78 (0.37 to 1.65)¶	Very low; downgraded for risk of bias, inconsistency and imprecision
Mobilization/rehabilitation interventions				
Mobility	3	330	0.29 (0.17 to 0.42)	Moderate; downgraded for risk of bias
Activities of daily living	1	182	0.48 (0.28 to 0.67)	Moderate; downgraded for risk of bias
Cognitive function	1	116	0.12 (-0.10 to 0.34)	Low; downgraded for risk of bias and imprecision
Falls	1	184	RR 0.88 (0.69 to 1.12)¶	Low; downgraded for risk of bias and imprecision

shown that older adults with frailty can improve mobility and ADL status with progressive resistance-based (muscle-strengthening) training.^{62,65,68,70-73}

We did not compare the effectiveness of different types or categories of physical activity in reducing frailty and frailty-related outcomes in older adults with frailty or pre-frailty. Previous reviews also showed that mixed physical

activity and muscle-strengthening interventions were the most common types of physical activity investigated in older adults with frailty.^{9,62,66,74,75} Other types of exercise, such as aerobic or endurance training, balance training, and flexibility or stretching training, have not been studied sufficiently, and their effectiveness has not been established.⁷⁴ We also did not analyze interventions based on frequency,

Table 4 (part 2 of 2): Physical activity intervention effect estimates for outcomes and certainty of evidence, for all interventions combined and by intervention category

Category; outcome*†	No. of studies	No. of participants	SMD‡ (95% CI)	GRADE rating
Mixed interventions				
Mobility	9	636	0.75 (0.40 to 1.10)	Moderate; downgraded for risk of bias
Activities of daily living	5	414	0.64 (0.004 to 1.27)	Low; downgraded for risk of bias and inconsistency
Cognitive function	2	180	0.62 (0.12 to 1.11)	Moderate; downgraded for risk of bias
Quality of life	5	430	0.68 (0.16 to 1.21)	Moderate; downgraded for risk of bias
Frailty	3	199	-1.57 (-2.57 to -0.57)	Moderate; downgraded for risk of bias
	3	1472	RR 0.72 (0.63 to 0.83)¶	Moderate; downgraded for risk of bias
Falls	1	82	-0.37 (-0.81 to 0.07)	Low; downgraded for risk of bias and imprecision
	1	98	RR 0.62 (0.16 to 2.47)¶	Low; downgraded for imprecision
Fatigue level	2		-0.23 (-0.85 to 0.39)	Very low; downgraded for risk of bias, inconsistency and imprecision
Use of health care services	1	82	-0.21 (-0.65 to 0.23)	Low; downgraded for risk of bias and imprecision
	1	98	RR 0.52 (0.05 to 5.56)¶	Low; downgraded for imprecision

Note: CI = confidence interval, GRADE = Grading of Recommendation, Assessment, Development and Evaluations,²² RR = risk ratio, SMD = standardized mean difference. *Mobility: gait speed, Timed Up and Go test, Sit-to-Stand test, balance test, Short Physical Performance Battery; cognitive function: Loewenstein Occupational Therapy Cognitive Assessment, Mini-Mental State Examination; falls: actual falls, fall rate, fall incidence; use of health care services: hospital admissions, emergency department visits.

†For aerobic interventions, there were no data for quality of life, frailty, falls or use of health care services; for muscle-strengthening interventions, there were no data for fatigue level or use of health care services; and for mobilization/rehabilitation interventions, there were no data for quality of life, frailty, fatigue level or use of health care services.

‡Except where noted otherwise.

§Large: -0.8; medium: -0.5; small: -0.2.

¶Dichotomous outcome.

intensity or duration, as this was not always clearly reported. Previous investigators have also struggled to analyze these data.^{9,15,65,74,76}

Physical activity interventions do not appear to introduce undue harm for older adults with frailty. Several studies reported no occurrences of adverse events, and minor events, such as aches, minor pains and fatigue, were managed successfully through adjustments made to the training protocols. For studies with more severe adverse events (e.g., falls, hospital admission), occurrence rates were similar between intervention and control groups,⁴⁴ and participant retention in these studies remained high (79%⁴⁴ to 92%³¹). Engaging in physical activity comes with inherent risks, but there are also well-known risks associated with sedentary time in older adults.⁷⁷ Although it is not realistic for all risk of harm to be eliminated during physical interventions, it is important that future studies and clinical and community programs follow published preparticipation screening recommendations to minimize risk to participants.⁷⁸ Importantly, as was evident from the high level of supervision in the studies in this review, these programs should engage professionals with the requisite skills and knowledge (such as nurses, physiotherapists and kinesiologists) to be able to adjust training protocols for this high-risk population as appropriate.

Research explicitly identifying prefrail and frail older adults, and consistency in the use of frailty identification tools are still lacking. In our review, very few studies measured frailty as a postintervention outcome. Likewise, the wide range of outcome measures used across studies, including unvalidated tools to assess frailty, may reflect the lack of clarity in frailty identification and diagnosis. Well-designed clinical trials that have explicit definitions of frailty and use outcomes that reflect frailty identification and diagnosis are needed to inform clear interventions that prevent or delay frailty progression in older adults.

We identified only 1 study specific to aerobic physical activity, so further research in this area is warranted.

Although the diversity of outcomes reflects the use of physical activity interventions in clinical and real-world settings, 54% of studies did not report the occurrence of adverse events. This limits our knowledge on the safety of the interventions and hampered comparison of studies.

Limitations

We used the inclusion criterion of frailty in an attempt to identify a more homogeneous population. However, it led to a large number of exclusions, which may have resulted in the exclusion of potentially frail populations or those with overlapping conditions. Furthermore, the variety of tools and




Figure 2: Effects of overall physical activity interventions on measures of mobility. Weights are from random-effects (RE) multilevel model analysis. Note: CI = confidence interval, df = degrees of freedom, Int. = intervention, SMD = standardized mean difference.

definitions used to describe participants still made for a diverse population of study participants that was subject to the authors' interpretation and description. We extracted outcomes immediately after the interventions, which leaves the long-lasting effects of the interventions unknown.

The practicality and uptake of exercises in real-world settings is unknown from this review. Dent and colleagues¹ noted that community-based programs for older adults with frailty often fall short of evidence-based recommendations, and that adherence to physical activity programs is poor among older adults for reasons such as fear of falling, lack

of self-belief, attitude, and adverse social and environmental influencers. Although we did not observe any significant asymmetry across funnel plots for publication bias, studies were small (< 300 participants) and had risk of bias concerns. Our search protocol had limitations, including restriction to English-language citations, our search end date of July 2019 and the fact that we did not search the grey literature.

Although there may be concerns about the reuse of the same participants from the same study to contribute data for multiple measures of a given outcome, our group agreed




Figure 3: Effects of overall physical activity interventions on measures of activities of daily living (A) and cognitive function (B). Weights are from random-effects (RE) multilevel model analysis. Note: ADL = activity of daily living, CI = confidence interval, df = degrees of freedom, GARS = Groningen Activity Restriction Scale, IADL = Instrumental Activities of Daily Living, Int. = intervention, LFFDI = Late-Life Function and Disability Instrument, LOTCA-G = Loewenstein Occupational Therapy Cognitive Assessment – Geriatric version, MMSE = Mini-Mental State Examination, RBANS = Repeatable Battery for the Assessment of Neuropsychological Status, SMD = standardized mean difference.

that this approach is preferred over selective reporting of effect sizes for a given outcome or averaging outcome measures from the same study to conduct a conventional 2-level meta-analysis. Both selective reporting (i.e., choosing

1 outcome measure for a given outcome) and averaging all outcome measures result in the potential loss and dilution of relevant information, and may produce misleading, inaccurate and biased results. These ad hoc approaches may




Figure 4: Effects of overall physical activity interventions on measures of quality of life (A) and frailty (B [continuous] and C [dichotomous]). Weights are from random-effects (RE) multilevel model analysis. Note: CI = confidence interval, df = degrees of freedom, DL = DerSimonian and Laird, EQ-5D = EuroQol-5 Dimension, RR = risk ratio, SF-36 = 36-Item Short Form Health Survey, SMD = standardized mean difference, SoF = Study of Osteoporotic Fractures criteria for frailty.

also lead to missed opportunities to use all available data to address the relevant research questions.⁷⁹

Conclusion

This review adds to the body of evidence identifying physical activities that benefit components of frailty in older adults, such as mobility, quality of life, cognitive function and ADLs, along with frailty itself. Our use of a strict inclusion criterion that attempted to identify a prefrail or frail population showed that studies need to better define frailty to ensure clear identification of older adults who would benefit from such interven-

tions. This would support the development of clear recommendations and facilitate the adoption and implementation of effective interventions into clinical practice.

References

- Dent E, Morley JE, Cruz-Jentoft AJ, et al. Physical frailty: ICFSR international clinical practice guidelines for identification and management. *J Nutr Health Aging* 2019;23:771-87.
- Muscledere J, Andrew MK, Bagshaw SM, et al. Screening for frailty in Canada's health care system: a time for action. *Can J Aging* 2016;35:281-97.
- Arantes PMM, Alencar MA, Dias RC, et al. Physical therapy treatment on frailty syndrome: systematic review. *Rev Bras Fisioter* 2009;13:365-75.

4. Puts MTE, Toubasi S, Andrew MK, et al. Interventions to prevent or reduce the level of frailty in community-dwelling older adults: a scoping review of the literature and international policies. *Age Ageing* 2017;46:383-92.
5. Negm AM, Kennedy CC, Thabane L, et al. Management of frailty: a systematic review and network meta-analysis of randomized controlled trials. *J Am Med Dir Assoc* 2019;20:1190-8.
6. Bock JO, König HH, Brenner H, et al. Associations of frailty with health care costs: results of the ESTHER cohort study. *BMC Health Serv Res* 2016;16:128.
7. Marcucci M, Damanti S, Germini F, et al. Interventions to prevent, delay or reverse frailty in older people: a journey towards clinical guidelines. *BMC Med* 2019;17:193.
8. Keyser JJ. Does late-life physical activity or exercise prevent or minimize disability? A critical review of the scientific evidence. *Am J Prev Med* 2003;25 (Suppl 2):129-36.
9. Paw MJMCA, van Uffelen JGZ, Riphagen I, et al. The functional effects of physical exercise training in frail older people: a systematic review. *Sports Med* 2008;38:781-93.
10. Chou CH, Hwang CL, Wu YT. Effect of exercise on physical function, daily living activities, and quality of life in the frail older adults: a meta-analysis. *Arch Phys Med Rehabil* 2012;93:237-44.
11. Kidd T, Mold F, Jones C, et al. What are the most effective interventions to improve physical performance in pre-frail and frail adults? A systematic review of randomised control trials. *BMC Geriatr* 2019;19:184.
12. DiPietro L, Buchner DM, Marquez DX, et al. New scientific basis for the 2018 U.S. Physical Activity Guidelines. *J Sport Health Sci* 2019;8:197-200.
13. Physical activity guidelines for older adults. London (UK): National Health Service; reviewed 2019 Oct. 8. Available: www.nhs.uk/live-well/exercise/physical-activity-guidelines-older-adults/ (accessed 2020 July 6).
14. Latham NK, Anderson CS, Lee A, et al. Randomised trial of home-based resistance training to improve physical health in frail elderly. *Intern Med J* 2003;33:A33.
15. Nash KCM. The effects of exercise on strength and physical performance in frail older people: a systematic review. *Rev Clin Gerontol* 2012;22:274-85.
16. Nowalk MP, Prendergast JM, Bayles CM, et al. A randomized trial of exercise programs among older individuals living in two long-term care facilities: the FallsFREE program. *J Am Geriatr Soc* 2001;49:859-65.
17. Racey MA, Ali MU, Sherifali D, et al.; Canadian Frailty Network. Effectiveness of nutrition interventions and combined nutrition and physical activity interventions in older adults with frailty or prefrailty: a systematic review and meta-analysis. *CMAJ Open* 2021;9:E744-56.
18. Moher D, Liberati A, Tetzlaff J, et al.; PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *J Clin Epidemiol* 2009;62:1006-12.
19. Higgins JPT, Li T, Deeks JJ. Choosing effect measures and computing estimates of effect. In: Higgins JPT, Thomas J, Chandler J, et al., editors. *Cochrane handbook for systematic reviews of interventions* version 6.1. London (UK): Cochrane; updated 2021:143-76.
20. Higgins JPT, Altman DG, Gøtzsche PC, et al.; Cochrane Statistical Methods Group. The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ* 2011;343:d5928.
21. Wells G, Shea B, O'Connell D, et al. The Newcastle-Ottawa Scale (NOS) for assessing the quality of non-randomized studies in meta-analyses. Ottawa: Ottawa Hospital Research Institute.
22. Schünemann H, Brożek J, Guyatt G, et al., editors. *GRADE handbook for grading quality of evidence and strength of recommendations*. London (UK): GRADE Working Group; updated 2013.
23. DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials* 1986;7:177-88.
24. Assink M, Wibbelink CJM. Fitting three-level meta-analytic models in R: a step-by-step tutorial. *Quant Methods Psychol* 2016;12:154-74.
25. Cohen J. *Statistical power analysis for the behavioral sciences*. 2nd ed. Mahwah (NJ): Lawrence Erlbaum Associates; 1988.
26. Pastor DA, Lazowski RA. On the multilevel nature of meta-analysis: a tutorial, comparison of software programs, and discussion of analytic choices. *Multivariate Behav Res* 2018;53:74-89.
27. Page MJ, Higgins JPT, Sterne JAC. Assessing risk of bias due to missing results in a synthesis. In: Higgins JPT, Thomas J, Chandler J, et al., editors. *Cochrane handbook for systematic reviews of interventions* version 6.2. London (UK): Cochrane; updated 2021:205-28.
28. Chen R, Wu Q, Wang D, et al. Effects of elastic band exercise on the frailty states in pre-frail elderly people. *Physiother Theory Pract* 2020;36:1000-8.
29. Losa-Reyna J, Baltasar-Fernandez I, Alcazar J, et al. Effect of a short multi-component exercise intervention focused on muscle power in frail and pre frail elderly: a pilot trial. *Exp Gerontol* 2019;115:114-21.
30. Liu Z, Hsu FC, Trombetti A, et al.; LIFE Study investigators. Effect of 24-month physical activity on cognitive frailty and the role of inflammation: the LIFE randomized clinical trial. *BMC Med* 2018;16:185.
31. Clegg A, Barber S, Young J, et al. The Home-based Older People's Exercise (HOPE) trial: a pilot randomised controlled trial of a home-based exercise intervention for older people with frailty. *Age Ageing* 2014;43:687-95.
32. Liu JWY, Lai CK, Siu PM, et al. An individualized exercise programme with and without behavioural change enhancement strategies for managing fatigue among frail older people: a quasi-experimental pilot study. *Clin Rehabil* 2017; 31:521-31.
33. Yoon DH, Lee JY, Song W. Effects of resistance exercise training on cognitive function and physical performance in cognitive frailty: a randomized controlled trial. *J Nutr Health Aging* 2018;22:944-51.
34. Takatori K, Matsumoto D, Nishida M, et al. Benefits of a novel concept of home-based exercise with the aim of preventing aspiration pneumonia and falls in frail older women: a pragmatic controlled trial. *BMJ Open Sport Exerc Med* 2016;2:e000127.
35. Tarazona-Santabalbina FJ, Gomez-Cabrera MC, Pérez-Ros P, et al. A multi-component exercise intervention that reverses frailty and improves cognition, emotion, and social networking in the community-dwelling frail elderly: a randomized clinical trial. *J Am Med Dir Assoc* 2016;17:426-33.
36. Ng TP, Feng L, Nyunt MSZ, et al. Nutritional, physical, cognitive, and combination interventions and frailty reversal among older adults: a randomized controlled trial. *Am J Med* 2015;128:1225-36.e1.
37. Kwon J, Yoshida Y, Yoshida H, et al. Effects of a combined physical training and nutrition intervention on physical performance and health-related quality of life in prefrail older women living in the community: a randomized controlled trial. *J Am Med Dir Assoc* 2015;16:263.e1-8.
38. Tieland M, Verdijk LB, de Groot LCPGM, et al. Handgrip strength does not represent an appropriate measure to evaluate changes in muscle strength during an exercise intervention program in frail older people. *Int J Sport Nutr Exerc Metab* 2015;25:27-36.
39. Giné-Garriga M, Guerra M, Unnithan VB. The effect of functional circuit training on self-reported fear of falling and health status in a group of physically frail older individuals: a randomized controlled trial. *Ageing Clin Exp Res* 2013;25:329-36.
40. Daniel K. Wii-hab for pre-frail older adults. *Rehabil Nurs* 2012;37:195-201.
41. Giné-Garriga M, Guerra M, Pagés E, et al. The effect of functional circuit training on physical frailty in frail older adults: a randomized controlled trial. *J Aging Phys Act* 2010;18:401-24.
42. Faber MJ, Bosscher RJ, Paw MJCA, et al. Effects of exercise programs on falls and mobility in frail and pre-frail older adults: a multicenter randomized controlled trial. *Arch Phys Med Rehabil* 2006;87:885-96.
43. Binder EF, Schechtman KB, Ehsani AA, et al. Effects of exercise training on frailty in community-dwelling older adults: results of a randomized, controlled trial. *J Am Geriatr Soc* 2002;50:1921-8.
44. Gill TM, Baker DI, Gottschalk M, et al. A program to prevent functional decline in physically frail, elderly persons who live at home. *N Engl J Med* 2002;347:1068-74.
45. de Jong N, Paw MJCA, de Graaf C, et al. Effect of dietary supplements and physical exercise on sensory perception, appetite, dietary intake and body weight in frail elderly subjects. *Br J Nutr* 2000;83:605-13.
46. Brown M, Sinacore DR, Ehsani AA, et al. Low-intensity exercise as a modifier of physical frailty in older adults. *Arch Phys Med Rehabil* 2000;81:960-5.
47. Yarasheski KE, Pak-Loduca J, Hasten DL, et al. Resistance exercise training increases mixed muscle protein synthesis rate in frail women and men ≥ 76 yr old. *Am J Physiol* 1999;277:E118-25.
48. Chandler JM, Duncan PW, Kochersberger G, et al. Is lower extremity strength gain associated with improvement in physical performance and disability in frail, community-dwelling elders? *Arch Phys Med Rehabil* 1998; 79:24-30.
49. Kuo MC, Chen CM, Jeng C. A randomized controlled trial of the prescribed stepper walking program in preventing frailty among the dwelling elderly: application of comprehensive geriatric assessment. *Top Geriatr Rehabil* 2018;34: 223-33.
50. Tsang HWH, Lee JLC, Au DWH, et al. Developing and testing the effectiveness of a novel health qigong for frail elders in Hong Kong: a preliminary study. *Evid Based Complement Alternat Med* 2013;2013:827392.
51. Chen KM, Chen MH, Lin MH, et al. Effects of yoga on sleep quality and depression in elders in assisted living facilities. *J Nurs Res* 2010;18:53-61.
52. Yamada M, Arai H. Self-management group exercise extends healthy life expectancy in frail community-dwelling older adults. *Int J Environ Res Public Health* 2017;14:531.
53. Yamada M, Arai H, Sonoda T, et al. Community-based exercise program is cost-effective by preventing care and disability in Japanese frail older adults. *J Am Med Dir Assoc* 2012;13:507-11.
54. Villareal DT, Steger-May K, Schechtman KB, et al. Effects of exercise training on bone mineral density in frail older women and men: a randomized controlled trial. *Age Ageing* 2004;33:309-12.
55. Ehsani AA, Spina RJ, Peterson LR, et al. Attenuation of cardiovascular adaptations to exercise in frail octogenarians. *J Appl Physiol (1985)* 2003;95:1781-8.
56. Binder EF, Yarasheski KE, Steger-May K, et al. Effects of progressive resistance training on body composition in frail older adults: results of a randomized, controlled trial. *J Gerontol A Biol Sci Med Sci* 2005;60:1425-31.
57. Paw MJCA, de Jong N, Schouten EG, et al. Physical exercise and/or enriched foods for functional improvement in frail, independently living elderly: a randomized controlled trial. *Arch Phys Med Rehabil* 2001;82:811-7.
58. Paw MJMCA, de Jong N, Schouten EG, et al. Physical exercise or micronutrient supplementation for the wellbeing of the frail elderly? A randomised controlled trial. *Br J Sports Med* 2002;36:126-31.

59. Gill TM, Baker DI, Gottschalk M, et al. A prehabilitation program for the prevention of functional decline: effect on higher-level physical function. *Arch Phys Med Rehabil* 2004;85:1043-9.
60. Ng TP, Ling LHA, Feng L, et al. Cognitive effects of multi-domain interventions among pre-frail and frail community-living older persons: randomized controlled trial. *J Gerontol A Biol Sci Med Sci* 2018;73:806-12.
61. Ng TP, Nyunt MSZ, Feng L, et al. Multi-domains lifestyle interventions reduces depressive symptoms among frail and pre-frail older persons: randomized controlled trial. *J Nutr Health Aging* 2017;21:918-26.
62. Lopez P, Pinto RS, Radaelli R, et al. Benefits of resistance training in physically frail elderly: a systematic review. *Aging Clin Exp Res* 2018;30:889-99.
63. Giné-Garriga M, Roqué-Fíguls M, Coll-Planas L, et al. Physical exercise interventions for improving performance-based measures of physical function in community-dwelling, frail older adults: a systematic review and meta-analysis. *Arch Phys Med Rehabil* 2014;95:753-69.e3.
64. Daniels R, van Rossum E, de Witte L, et al. Interventions to prevent disability in frail community-dwelling elderly: a systematic review. *BMC Health Serv Res* 2008;8:278.
65. Theou O, Stathokostas L, Roland KP, et al. The effectiveness of exercise interventions for the management of frailty: a systematic review. *J Aging Res* 2011;2011:569194.
66. Cadore EL, Rodríguez-Mañas L, Sinclair A, et al. Effects of different exercise interventions on risk of falls, gait ability, and balance in physically frail older adults: a systematic review. *Rejuvenation Res* 2013;16:105-14.
67. de Vries NM, Staal JB, van der Wees PJ, et al. Patient-centred physical therapy is (cost-) effective in increasing physical activity and reducing frailty in older adults with mobility problems: a randomized controlled trial with 6 months follow-up. *J Cachexia Sarcopenia Muscle* 2016;7:422-35.
68. de Labra C, Guimaraes-Pinheiro C, Maseda A, et al. Effects of physical exercise interventions in frail older adults: a systematic review of randomized controlled trials. *BMC Geriatr* 2015;15:154.
69. Macdonald SHF, Travers J, Shé ÉN, et al. Primary care interventions to address physical frailty among community-dwelling adults aged 60 years or older: a meta-analysis. *PLoS One* 2020;15:e0228821.
70. Apóstolo J, Cooke R, Bobrowicz-Campos E, et al. Effectiveness of interventions to prevent pre-frailty and frailty progression in older adults: a systematic review. *JBI Database System Rev Implement Rep* 2018;16:140-232.
71. Bauman A, Merom D, Bull FC, et al. Updating the evidence for physical activity: summative reviews of the epidemiological evidence, prevalence, and interventions to promote "active aging." *Gerontologist* 2016;56(Suppl 2):S268-80.
72. Mundell BF, Kremers HM, Visscher S, et al. Predictors of receiving a prosthesis for adults with above-knee amputations in a well-defined population. *PM R* 2016;8:730-7.
73. Raymond MJ, Bramley-Tzeretos RE, Jeffs KJ, et al. Systematic review of high-intensity progressive resistance strength training of the lower limb compared with other intensities of strength training in older adults. *Arch Phys Med Rehabil* 2013;94:1458-72.
74. Jadcak AD, Makwana N, Luscombe-Marsh N, et al. Effectiveness of exercise interventions on physical function in community-dwelling frail older people: an umbrella review of systematic reviews. *JBI Database System Rev Implement Rep* 2018;16:752-75.
75. Sherrington C, Fairhall NJ, Wallbank GK, et al. Exercise for preventing falls in older people living in the community. *Cochrane Database Syst Rev* 2019;(1):CD012424.
76. Bray NW, Smart RR, Jakobi JM, et al. Exercise prescription to reverse frailty. *Appl Physiol Nutr Metab* 2016;41:1112-6.
77. Copeland JL, Ashe MC, Biddle SJ, et al. Sedentary time in older adults: a critical review of measurement, associations with health, and interventions. *Br J Sports Med* 2017;51:1539.
78. Riebe D, Franklin BA, Thompson PD, et al. Updating ACSM's recommendations for exercise preparticipation health screening. *Med Sci Sports Exerc* 2015;47:2473-9.
79. Cheung MWL. A guide to conducting a meta-analysis with non-independent effect sizes. *Neuropsychol Rev* 2019;29:387-96.

Affiliations: McMaster Evidence Review and Synthesis Team (Racey, Ali, Sherifali, Fitzpatrick-Lewis, Lewis, Jovkovic); School of Nursing (Racey, Sherifali, Fitzpatrick-Lewis), Department of Clinical Epidemiology & Biostatistics (Ali) and School of Rehabilitation Science (Tang), Faculty of Health Sciences, McMaster University, Hamilton, Ont.; Faculty of Kinesiology (Bouchard), University of New Brunswick; Cardiometabolic Exercise

& Lifestyle Laboratory (Bouchard), Fredericton, NB; Department of Family and Emergency Medicine (Giguère), Laval University, Québec, Que.; Cumming School of Medicine (Holroyd-Leduc), University of Calgary, Calgary, Alta.; University of Alberta (Gramlich); Nutrition Services (Gramlich), Alberta Health Services, Edmonton, Alta.; Schlegel-University of Waterloo Research Institute for Aging (Keller) and Department of Kinesiology (Keller), University of Waterloo, Waterloo, Ont.; Canadian Frailty Network (Prorok, Kim, Lorbergs, Muscedere), Kingston, Ont.

Contributors: Megan Racey, Muhammad Usman Ali, Diana Sherifali, Donna Fitzpatrick-Lewis, Leah Gramlich, Heather Keller, Jayna Holroyd-Leduc, Anik Giguère, Ada Tang, Danielle Bouchard, Jeanette Prorok, Perry Kim, Amanda Lorbergs and John Muscedere designed the study. Megan Racey, Muhammad Usman Ali, Diana Sherifali, Donna Fitzpatrick-Lewis, Ruth Lewis and Milos Jovkovic acquired and analyzed the data and drafted the manuscript. Megan Racey, Muhammad Usman Ali, Diana Sherifali and Donna Fitzpatrick-Lewis interpreted the data. Leah Gramlich, Heather Keller, Jayna Holroyd-Leduc, Anik Giguère, Ada Tang, Danielle Bouchard, Jeanette Prorok, Perry Kim, Amanda Lorbergs and John Muscedere revised the manuscript critically for important intellectual content. All of the authors approved the final version to be published and agreed to be accountable for all aspects of the work.

Members of the Canadian Frailty Network: John Muscedere (Scientific Director), Carol Barrie (Executive Director), Perry Kim (Assistant Scientific Director), Amanda Lorbergs (Manager, Research and Knowledge Translation), Jeanette Prorok (Manager, Special Projects and Initiatives), Kyle Plumb (Project Manager), Mat LaBranche (Finance & Administration), Amy Doyle (Manager, Strategic Partnership Development and Communications), Kelsey MacIntosh (Health Promotion Coordinator)

Funding: This research was funded by the McMaster Institute for Research and Aging, the McMaster Evidence Review and Synthesis Team (MERST) and the Canadian Frailty Network (CFN) (Technology Evaluation in the Elderly Network), which is supported by the Government of Canada through the Networks of Centres of Excellence program. The CFN was responsible for the conception of the research question and supported the editing of the manuscript. MERST was independently responsible for data extraction, analysis, interpretation and reporting. Together, these funds supported the work of Megan Racey as a postdoctoral fellow. Diana Sherifali holds the Heather M. Arthur Population Health Research Institute/Hamilton Health Sciences Chair in Inter-professional Health Research, which supported her role in this work.

Disclaimer: No one from the Canadian Frailty Network had any involvement in the data extraction or analysis for the review. All screening of articles, data extraction, and data analysis and interpretation were conducted by the first 6 authors, independent from the steering committee and funders.

Content licence: This is an Open Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY-NC-ND 4.0) licence, which permits use, distribution and reproduction in any medium, provided that the original publication is properly cited, the use is noncommercial (i.e., research or educational use), and no modifications or adaptations are made. See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

Data sharing: Data used in this review can be requested for confidential research purposes by contacting the corresponding author(s).

Acknowledgement: The authors thank Angela Eady for developing the search strategy.

Supplemental information: For reviewer comments and the original submission of this manuscript, please see www.cmajopen.ca/content/9/3/E728/suppl/DC1.