

Appendix 2 (as supplied by the authors): Univariable quantile regression models predicting diagnostic intervals for patients first presenting in the ER (in days)

			Median	95% CI	p
			difference		
	Intercept		3.848	-10.99 - 18.69	0.6094
Age	(per 10 years)		0.494	-1.57 - 2.56	0.6369
	Intercept		6.000	2.40 - 9.60	0.0012
Stage	I/II		16.000	-4.59 - 36.59	0.1270
	III/IV		(reference)		
	Unknown		0.000	-5.70 - 5.70	1.0000
	Intercept		6.000	2.07 - 9.93	0.0030
Morphology	Serous Carcinoma		0.000	-6.55 - 6.55	1.0000
	Unclassified Epithelial		(reference)		
	Clear Cell / Endometrioid		25.000	-3.47 - 53.47	0.0849
	Mucinous		9.000	-16.30 - 34.30	0.4835
	Other		-2.000	-9.53 - 5.53	0.6006
	*other epithelial-stromal, and miscellaneous and unspecified				
	Intercept		10.000	-0.63 - 20.63	0.0651
Residence	Winnipeg		-3.000	-14.14 - 8.14	0.5955
	Non-Winnipeg		(reference)		
	Intercept		7.000	3.83 - 10.17	<.0001
Period of diagnosis	2008 and later		1.000	-4.52 - 6.52	0.7211
	2007 and earlier		(reference)		
Income	<i>Contrasts</i>				
	R4-R5 vs R1-R3		-7.958	-31.25 - 15.34	0.5009
	U4-U5 vs U1-U3		-3.000	-8.92 - 2.92	0.3188

Appendix to: Love AJ, Lambert P, Turner D, et al. Diagnostic and referral intervals for Manitoba women with epithelial ovarian cancer — the Manitoba Ovarian Cancer Outcomes (MOCO) study group: a retrospective cross-sectional study. *CMAJ Open* 2017.

DOI:10.9778/cmajo.20160100. Copyright © 2017 The Author(s) or their employer(s).

To receive this resource in an accessible format, please contact us at cmajgroup@cmaj.ca.

*R=rural; U=urban; 1=poorest;5=richest					
	Intercept		6.000	2.73 - 9.27	0.0004
Comorbidities	High/Very High		-3.000	-2.70 - 8.70	0.2999
(Resource utilization	Moderate and lower		(reference)		
band)					
	Intercept		10.042	3.45 - 16.63	0.0030
Abdominal distension	Yes		-5.042	-12.34 - 2.26	0.1747
	No		(reference)		
	Intercept		7.000	3.78 - 10.22	<.0001
Incidental	Yes		6.000	-101.63 - 113.63	0.9125
	No		(reference)		
	Intercept		8.000	4.13 - 11.87	0.0001
Bowel symptoms	Yes		-4.000	-9.44 - 1.44	0.1485
	No		(reference)		
	Intercept		8.000	4.41 - 11.59	<.0001
Nausea	Yes		-5.000	-9.04 - -0.96	0.0156
	No		(reference)		
	Intercept		8.000	4.56 - 11.44	<.0001
Decreased appetite	Yes		-6.000	-10.20 - -1.80	0.0054
	No		(reference)		
	Intercept		8.000	4.27 - 11.73	<.0001
Respiratory symptoms	Yes		-3.000	-9.57 - 3.57	0.3686
	No		(reference)		

	Intercept		8.000	4.96 - 11.04	<.0001
Weight change	Yes		-6.000	-11.95 - -0.05	0.0480
	No		(reference)		
	Intercept		7.000	4.13 - 9.87	<.0001
Urinary symptoms	Yes		17.000	-26.08 - 60.08	0.4370
	No		(reference)		
	Intercept		-		
Abnormal bleeding	Yes		-		
	No		-		
	Intercept		-		
Postmenopausal	Yes		-		
bleeding	No		-		
	Intercept		7.958	5.04 - 10.87	<.0001
Palpable mass	Yes		-4.958	-16.20 - 6.28	0.3852
	No		(reference)		
	Intercept		8.000	4.75 - 11.25	<.0001
Weakness	Yes		-5.000	-10.75 - 0.75	0.0878
	No		(reference)		
	Intercept		4.000	1.82 - 6.18	0.0004
Abdominal pain and vomiting	Pain and no emesis		7.000	2.26 - 11.74	0.0040
and emesis	Pain and emesis		-2.000	-4.69 - 0.69	0.1434
	No pain or emesis		(reference)		