

[Type text]

Appendix 5 (as supplied by the authors): Univariable quantile regression models predicting referral intervals for patients first presenting outside the ER (in days)

		Median	95% CI	p
		difference		
Intercept		64.563	33.26 - 95.87	0.0001
Age	(per 10 years)	-1.565	-6.43 - 3.29	0.5270
Intercept		43.000	35.82 - 50.18	<.0001
Stage	I/II	18.000	6.96 - 29.04	0.0015
	III/IV	(reference)		
	Unknown	32.958	6.05 - 59.87	0.0165
Intercept		53.000	38.64 - 67.36	<.0001
Morphology	Serous Carcinoma	-3.958	-20.74 - 12.82	0.6431
	Unclassified Epithelial	(reference)		
	Clear Cell / Endometrioid	7.000	-16.97 - 30.97	0.5662
	Mucinous	17.042	-6.28 - 40.36	0.1517
	Other	3.042	-23.50 - 29.58	0.8218
	*other epithelial-stromal, and miscellaneous and unspecified			
Intercept		54.958	45.47 - 64.45	<.0001
Residence	Winnipeg	0.000	-12.10 - 12.10	1.0000
	Non-Winnipeg	(reference)		
Intercept		49.042	41.09 - 56.99	<.0001
Period of diagnosis	2008 and later	11.958	-1.73 - 25.64	0.0866
	2007 and earlier	(reference)		
Income	<i>Contrasts</i>			
	R4-R5 vs R1-R3	-3.000	-21.83 - 15.83	0.7526
	U4-U5 vs U1-U3	-1.000	-16.04 - 14.04	0.8960

Appendix to: Love AJ, Lambert P, Turner D, et al. Diagnostic and referral intervals for Manitoba women with epithelial ovarian cancer — the Manitoba Ovarian Cancer Outcomes (MOCO) study group: a retrospective cross-sectional study. *CMAJ Open* 2017. DOI:10.9778/cmajo.20160100. Copyright © 2017 The Author(s) or their employer(s).

To receive this resource in an accessible format, please contact us at cmajgroup@cmaj.ca.

[Type text]

*R=rural; U=urban; 1=poorest;5=richest				
Intercept		59.000	53.03 - 64.97	<.0001
Comorbidities	High/Very High	-15.000	-25.24 - -4.76	0.0042
(Resource utilization band)	Moderate and lower	(reference)		
Intercept		56.000	49.17 - 62.83	<.0001
Abdominal pain	Yes	-3.000	-14.78 - 8.78	0.6169
	No	(reference)		
Intercept		62.000	54.59 - 69.41	<.0001
Abdominal distension	Yes	-28.000	-39.44 - -16.56	<.0001
	No	(reference)		
Intercept		53.000	46.23 - 59.77	<.0001
Incidental	Yes	13.958	-6.29 - 34.20	0.1761
	No	(reference)		
Intercept		54.958	48.87 - 61.05	<.0001
Bowel symptoms	Yes	0.042	-21.01 - 21.09	0.9969
	No	(reference)		
Intercept		54.958	49.30 - 60.62	<.0001
Nausea	Yes	-7.958	-28.14 - 12.22	0.4387
	No	(reference)		
Intercept		56.000	49.99 - 62.01	<.0001
Decreased appetite	Yes	-14.000	-25.00 - -3.00	0.0127
	No	(reference)		

[Type text]

Intercept			54.958	48.80 - 61.12	<.0001
Respiratory symptoms	Yes		2.000	-28.39 - 32.39	0.8971
	No		(reference)		
Intercept			55.000	49.33 - 60.67	<.0001
Weight change	Yes		-9.000	-19.70 - 1.70	0.0990
	No		(reference)		
Intercept			55.000	49.48 - 60.52	<.0001
Urinary symptoms	Yes		-14.000	-48.14 - 20.14	0.4207
	No		(reference)		
Intercept			54.958	50.02 - 59.90	<.0001
Abnormal bleeding	Yes		7.083	-29.86 - 44.02	0.7064
	No		(reference)		
Intercept			53.000	46.42 - 59.58	<.0001
Postmenopausal bleeding	Yes		41.958	-11.13 - 95.05	0.1210
	No		(reference)		
Intercept			55.000	49.13 - 60.87	<.0001
Palpable mass	Yes		-2.000	-15.96 - 11.96	0.7783
	No		(reference)		
Intercept			55.000	49.21 - 60.79	<.0001
Weakness	Yes		-11.000	-26.94 - 4.94	0.1757
	No		(reference)		
Intercept			54.958	48.51 - 61.40	<.0001
Vomiting	Yes		-3.958	-30.28 - 22.37	0.7677

[Type text]

	No		(reference)		
--	----	--	-------------	--	--