

Appendix 1 (as supplied by the authors): Change in scores on Physician Readiness to Manage Intimate Partner Violence Survey subscales at baseline, immediately after, 3 and 12 months post-training*

Subscales**	Baseline mean (SD)	Immediately post-training mean (SD) [1]	3-months mean (SD) [1]	12-months mean (SD)	Multivariable linear regression model comparing baseline to 12-month scores
					Mean difference (95% CI)
Actual knowledge	26.59 (4.92)	30.06 (3.95)	29.04 (3.89)	29.09 (4.66)	2.50 (1.69, 3.32)
Perceived preparation	2.60 (1.08)	4.74 (1.14)	4.59 (1.12)	4.66 (1.15)	2.06 (1.88, 2.24)
Perceived knowledge	2.70 (1.12)	4.89 (1.01)	4.77 (1.05)	4.84 (1.05)	2.14 (1.96, 2.31)
Opinion subscale					
Preparation	3.73 (1.25)	4.75 (0.94)	4.74 (0.97)	4.83 (0.90)	1.10 (0.94, 1.26)
Legal requirements	3.38 (1.52)	5.10 (1.17)	4.91 (1.12)	4.94 (1.16)	1.57 (1.36, 1.78)
Workplace issues†	3.04 (0.95)	4.12 (0.82)	4.11 (0.88)	4.24 (0.88)	1.19 (1.04, 1.35)
Self-efficacy	3.56 (0.46)	4.12 (0.49)	4.09 (0.57)	4.12 (0.61)	0.56 (0.46, 0.67)
Alcohol and drugs	4.22 (0.57)	4.44 (0.58)	4.28 (0.46)	4.34 (0.52)	0.11 (0.02, 0.21)
Victim understanding	4.97 (0.70)	5.08 (0.77)	4.95 (0.78)	4.92 (0.70)	-0.05 (-0.16, 0.07)
Practice issues	5.81 (6.46)	9.62 (5.91)	11.83 (7.74)	11.94 (7.26)	6.12 (4.97, 7.27)

Note: CI=confidence interval, SD=standard deviation

* At 12 months, N=108. One participant missed all questions that form the workplace issues domain.

** Ranges and estimated minimally important clinical differences (MCIDs)[†] for subscales (range, MCID): **actual knowledge** (0 to 38, 2.42); **perceived preparation to manage IPV** (1 to 7, 0.55); **perceived knowledge of important IPV issues** (1 to 7, 0.55); **practice issues** (0 to 58, 3.06); **preparation** (1 to 7, 0.59); **legal requirements** (1 to 7, 0.77); **workplace issues** (1 to 7, 0.45); **self-efficacy** (1 to 7, 0.22); **alcohol/drugs** (1 to 7, 0.28); and **victim understanding** (1 to 7, 0.35). † MCIDs were estimated as half the subscales' standard deviations [2]

References

1. EDUCATE Investigators. Novel educational program improves readiness to manage intimate partner violence within the fracture clinic: a pretest-posttest study. *CMAJ Open*. 2018 Dec;6(4):E628–E636.
2. Norman GR, Sloan JA, Wyrwich KW. Interpretation of changes in health-related quality of life: the remarkable universality of half a standard deviation. *Med Care* 2003;41:582-92.