

1 **Supplemental Table 1.** Region of travel of travellers and country of origin of migrants in those
 2 presenting with Old world and New world CL.

Region/country	Total, n (%)	OW CL, n (%)	NW CL, n (%)
North America	37 (77.1)	9 (60.0)	28 (84.8)
Canada	35 (72.9)	9 (60.0)	26 (78.8)
United States	1 (2.1)	0 (0.0)	1 (3.0)
Mexico	1 (2.1)	0 (0.0)	1 (3.0)
Central America/Caribbean	2 (4.2)	0 (0.0)	2 (6.1)
Belize	1 (2.1)	0 (0.0)	1 (3.0)
Haiti	1 (2.1)	0 (0.0)	1 (3.0)
South America	2 (4.2)	0 (0.0)	2 (6.1)
Colombia	1 (2.1)	0 (0.0)	1 (3.0)
Peru	1 (2.1)	0 (0.0)	1 (3.0)
Europe	1 (2.1)	0 (0.0)	1 (3.0)
France	1 (2.1)	0 (0.0)	1 (3.0)
Middle East/South Central India	5 (10.4)	5 (33.3)	0 (0.0)
Syria	2 (4.2)	2 (13.3)	0 (0.0)
Iran	1 (2.1)	1 (6.7)	0 (0.0)
Afghanistan	2 (4.2)	2 (13.3)	0 (0.0)
North Africa	1 (2.1)	1 (6.7)	0 (0.0)
Algeria	1 (2.1)	1 (6.7)	0 (0.0)

3 OW CL: Old World Cutaneous Leishmaniasis; NW CL: New World Cutaneous Leishmaniasis

4
 5
 6
 7

8

9 **Supplemental Table 2.** Countries of exposure to CL

Country	Total (n, %) n=48	OW CL (n, %) n=15	NW (n, %) n=31*
Costa Rica	11 (24%)	0 (0%)	11 (35%)
Mexico	7 (15%)	0 (0%)	7 (23%)
Burkina Faso	3 (7%)	3 (20%)	0 (0%)
Belize	2 (4%)	0 (0%)	2 (6%)
Bolivia	2 (4%)	0 (0%)	2 (6%)
Colombia	2 (4%)	0 (0%)	2 (6%)
Morocco	2 (4%)	2 (13%)	0 (0%)
Peru	2 (4%)	0 (0%)	2 (6%)
Syria	2 (4%)	2 (13%)	0 (0%)
Tunisia	2 (4%)	2 (13%)	0 (0%)
Brazil	1 (2%)	0 (0%)	1 (3%)
China	1 (2%)	1 (7%)	0 (0%)
Ecuador	1 (2%)	0 (0%)	1 (3%)
French Guinea	1 (2%)	0 (0%)	1 (3%)
Guatemala	1 (2%)	0 (0%)	1 (3%)
Iran	1 (2%)	1 (7%)	0 (0%)
Israël	1 (2%)	1 (7%)	0 (0%)
Pakistan	1 (2%)	1 (7%)	0 (0%)
Panama	1 (2%)	0 (0%)	1 (3%)

10 OW CL: Old World Cutaneous Leishmaniasis; NW CL: New World Cutaneous Leishmaniasis

11 *There were 2 patients in the NW CL group where the country of exposure was not ascertainable.

12

13

14

15 **Supplemental Table 3. *Leishmania* species and country of exposure**

<i>Leishmania</i> spp.	Travellers and migrants (n=43)*	Top countries
New World n (%)	33	Costa Rica (11), Mexico (6), Bolivie (2), Colombie (2)
<i>Leishmania (V.) panamensis</i>	23	Costa Rica (11), Bolivie (2), Colombie (2)
<i>Leishmania (V.) braziliensis</i>	3	French Guinea (1), Belize (1), Peru (1)
<i>Leishmania (L.) mexicana</i>	7	Mexico (6), Belize (1)
Old World n (%)	10	Tunisia (2), Morocco (2)
<i>Leishmania (L.) major</i>	7	Tunisia (2), Morocco (2), Algeria (1)
<i>Leishmania (L.) tropica</i>	3	Pakistan (1), Israel (1), Syria (1)

16 *Speciation was available for 43 out of 48 travellers and migrants

17