## **Amoxicillin Allergy Survey**

	Birth: Card Number: Physician:	PATIENT STICK	ER		Date of Allergi	c Re	eaction:
are rep prescril serious amoxic followir	illin is a penicillin-bas orted in as many as bing amoxicillin, but , or anaphylactic, ro illin can be used saf	5 to 10% of pat many of these a eactions should fely when there h	ients. Often re not true a not use amo as previously	a realler oxici	eaction will lead patie gic reactions. Patie llin. There is grow en a mild, moderate	ents ents /ing e, or	se reactions to penicillins and physicians to avoid who have experienced research to show that delayed reaction. The e future risks associated
Sympt	oms:						
Please indicate if any of the following symptoms were part of your allergic reaction:  □ Loss of consciousness, fainting, or near fainting □ Difficulty breathing, wheezing □ Abdominal pain □ Swelling of the lips, tongue, or face □ Anaphylaxis							
If you selected one of the above answers, please STOP HERE. If you did not, then continue below:							
	ind of symptoms did y Rash Other		Itching				Blistering
	lid the symptoms star Within minutes or ho		Within 24h	rs			After 24hrs
How lo	ng did the symptoms l Hours	last:	Days				Weeks
Other reactions or allergies: Have you had an allergic reaction to another medication(s)?							
Has one of your parents had an allergic reaction to amoxicillin?							
Assurance of Accuracy: By signing below, you certify that the information provided in this survey is accurate and complete to best of knowledge. Withholding or providing misinformation will affect your physician's ability to treat you safely, and could lead to very dangerous medical complications.							
Patient	/Parent Name:						
Patient/Parent Signature:							
Office Use Only: □ Immediate □ Ig-E □ Severe □ Challenge							